

# Labcorp

Specimen Number 035		Patient ID [REDACTED]		Control Number [REDACTED]	Account Number [REDACTED]	Account Phone Number [REDACTED]	Route [REDACTED]
Patient Last Name [REDACTED]				Clinic [REDACTED]			
Patient First Name [REDACTED]		Patient Middle Name [REDACTED]		Accession Number [REDACTED]		Req/Ctrl#(ACC) [REDACTED]	
Patient SS# XXX-XX-XXXX	Patient Phone [REDACTED]	Total Volume [REDACTED]		Status Final	Date [REDACTED]	User [REDACTED]	
Age(Y/M/D) 27 / 3 / 109	Date of Birth [REDACTED]	Sex M	Fasting N	Physician Name [REDACTED]		NPI [REDACTED]	
Date and Time Collected 02/04/2015 18:55:00	Date Entered 02/05/2015 06:11:00	Date and Time Reported 02/05/2015 12:21:00		Patient Address [REDACTED]			

## Additional Information

### Test Ordered

RUBEOLA ANTIBODIES, IGG						
TEST	RESULT	FLAG	UNITS	REFERENCE INTERVAL	LAB	TECH
RUBEOLA AB, IGG	<25.0	Below Normal Low	AU/ML	IMMUNE >29.9	PDLCA	
<b>Comments</b> Negative <25.0 Equivocal 25.0 - 29.9 Positive >29.9 Presence of antibodies to Rubella is presumptive evidence of immunity except when acute infection is suspected.						

### Place Of service

[REDACTED]	[REDACTED]	[REDACTED]
------------	------------	------------

Patient:

DOB:

Control Number:

## Patient

██████████  
Male, ██████████  
Home: ██████████  
Mobile: ██████████  
Email: ██████████

Fasting: No

## Physician

██████████  
Genesis Healing Centers  
██████████  
F: ██████████  
Email: ██████████  
██████████

## Visit

Date: 2/14/2015

Time: 10:32 AM

Acc #: ██████████

## Test: Rubella Antibody, IgG

Result Name	Result value	Units	Flag	Reference Range
Measles virus Ab.IgG	27.4	AU/mL	EQUIVOCAL	

## Interpretative Note:

### Index Result Interpretation

#### < 25 = Negative

- Absence of detectable measles virus IgG antibodies.
- A negative result generally indicates that the patient has not been previously infected and is therefore susceptible to measles.
- If the subject has no history of measles has not been previously vaccinated and exposure to measles virus is nonetheless suspected despite a negative finding a second sample should be collected and tested no less than one to two weeks later.

#### 25 <= x <= 30 = Equivocal

- Equivocal samples should be retested. If the result remains equivocal after repeat testing a second sample should be collected no less than one to two weeks later.

#### > 30 = Positive

- Presence of detectable measles virus IgG antibodies. A positive result generally indicates past exposure to measles virus or previous vaccination.
- The magnitude of the measured result above the cutoff is not indicative of the amount of antibody present.
- Diagnosis of infectious diseases should not be established on the basis of a single test result but should be determined in conjunction with previous infection history clinical findings and other diagnostic procedures as well as in association with medical judgment.

Verified by repeat run.

## Performing Site:

Newark - 7373 Gateway Boulevard, Newark CA 94560